

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

BUNDESREPUBLIK DEUTSCHLAND

EP 00 / 1368



Bescheinigung

09/913631

Die Theragene Biomedical Laboratories GmbH in Martinsried/Deutschland hat eine Patentanmeldung unter der Bezeichnung

"Hormone-Hormone Receptor Complexes and Nucleic Acid Constructs and Their Use in Gene Therapy"

am 19. Februar 1999 beim Deutschen Patent- und Markenamt eingereicht.

Die angehefteten Stücke sind eine richtige und genaue Wiedergabe der ursprünglichen Unterlagen dieser Patentanmeldung.

Die Anmeldung hat im Deutschen Patent- und Markenamt vorläufig die Symbole C 07 H, C 12 N und A 61 K der Internationalen Patentklassifikation erhalten.

München, den 20. März 2000

Deutsches Patent- und Markenamt

Der Präsident

Im Auftrag

Dzierzon

Aktenzeichen: 199 07 099.7

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Abstract of the Disclosure

Belegexemplar
Darf nicht geändert werden



The invention provides a composition of matter comprising a nucleotide sequence comprising at least one hormone responsive element (HRE), said hormone responsive element being coupled to a hormone-hormone receptor complex and further provides nucleic acid constructs comprising at least one hormone responsive element (HRE), wherein the hormone responsive element regulates a gene encoding a human blood-clotting factor, the invention having applications in gene therapy, and particularly, in the treatment of human blood clotting disorders, such as hemophilia.

Hormone -Hormone Receptor Complexes and Nucleic Acid Constructs **and Their Use in Gene Therapy**

Background of the Invention

5

1. Object of the Invention

The invention provides a composition of matter comprising a nucleotide sequence comprising at least one hormone responsive element (HRE), said hormone responsive element being coupled to a hormone-hormone receptor complex. It further relates to nucleic acid constructs comprising at least one hormone responsive element and vectors comprising such constructs, wherein the hormone responsive element regulates a gene encoding a human blood-clotting factor. The nucleic acid constructs, plasmids, and compositions of matter of the invention have applications in gene therapy, particularly in the treatment of human blood clotting disorders, such as hemophilia. They may also be used to up- or down-regulate target genes and for the delivery of vaccines.

2. Summary of the Related Art

Gene therapy is a method that holds great promise for many diseases and disorders. In general, it involves the transfer of recombinant genes or transgenes into somatic cells to replace proteins with a genetic defect or to interfere with the pathological process of an illness. In principle, gene therapy is a simple method. In practice, many disadvantages must still be overcome.

Research in gene therapy has concentrated on ways to most effectively incorporate DNA into cells of a patient. Viral vectors are currently the widely used vehicles in clinical gene therapy approaches. In terms of efficacy in gene expression, the

viral delivery systems have major advantages over techniques using DNA-lipid formulations as delivery vehicles or over mechanical methods, such as the gene gun. Although there are a variety of viral systems tested for gene therapeutical strategies, retroviral vectors and adenoviral vectors are presently the most widely used vehicles (Salmons, B. and Gunzburg, W. H., *Hum. Gene Ther.*, Vol. 4, 129, 1993; Kasahara, N. A., et al., *Science*, Vol. 266, 1373, 1994; Ali, M., et al., *Gene Ther.*, Vol. 1, 367, 1994.). Still, these systems have major disadvantages, such as potential viral contamination. Other safety concerns continue to hamper the development of clinical application of gene therapy using these viral systems. For example, recombinant retroviruses have the disadvantage of random chromosomal integration, which may lead to activation of oncogenes or inactivation of tumor-suppressor genes. Also, repetitive use of recombinant adenoviruses has caused severe immunological problems (Elkon, K. B. et al., *Proc. Natl. Acad. Sci. USA*, Vol. 94, 9814, 1997). The humoral response resulted in the production of antibodies to adenovirus proteins preventing subsequent infection. Immunosuppressive drugs may ameliorate these effects, but they place an additional burden on the patient (Dai, Y., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 92, 1401, 1995).

Yet another viral delivery system involves adenoassociated virus (AAV). The AAV requires coinfection with an unrelated helper virus. Although such recombinant AAV virions have proven useful for introducing several small gene sequences into host cells, gene delivery systems based on those particles are limited by the relative small size of AAV particles. This feature greatly reduces the range of appropriate gene protocols. Moreover, the need to also use a helper virus adds a complicating factor to this delivery system (Muzyczka, N., *Curr. Top. Microbiol. Immunol.*, Vol. 158, 97, 1992).

Though safer, non-viral gene therapy approaches are also unsatisfactory. Problems with inefficient gene delivery or poor sustained expression are major drawbacks. Yet methods such as the direct injection of DNA into cellular compartments to mixtures of DNA with cationic lipids or polylysine allowing the transgene to cross the cell membrane more easily, have not overcome these hurdles (Felgner, P., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 84, 7413, 1987; Behr, J.-P., *Bioconjugate Chemistry*, Vol. 5, 382, 1994).

10

20

30

exogenous supply of Factor VIII to a patient still requires repeated doses throughout the lifetime of the patient, an inconvenient and expensive solution. Other forms of hemophilia include Hemophilia B, caused by a defect in the gene coding for Factor IX. The gene therapy systems described above have been attempted for the treatment of Hemophilia A and B with Factors VIII and IX, respectively. (See e.g., WO 94/29471). However, these systems have the disadvantages already discussed above.

The classical model of the action of hormones is based on the concept of binding interaction of the hormone to an intracellular receptor, located in the cytoplasm or the nucleus (Evans, R., *Science*, Vol. 240, 889, 1988). These intracellular receptors remain latent until exposed to their target hormone. When so exposed, the hormone receptor changes its conformation after the hormone is bound and translocates in the activated form into the cell nucleus where it binds as a dimer to hormone-responsive-elements (HRE) in the promoter region of hormone-regulated genes (Beato, M., *Cell*, Vol. 56, 335, 1989; O'Malley, B., et al., *Biol. Reprod.*, Vol. 46, 163, 1992). The HRE are enhancer elements usually located in the 5' flanking region of the specific hormone-induced gene.

An example of such intracellular receptors is the steroid receptor. Steroid receptors belong to a superfamily of ligand-dependent transcription factors characterized by a unique molecular structure. The centrally located highly conserved DNA-binding domain defines this superfamily. The second important and relatively invariant region is the COOH-terminal ligand-binding domain. An example of such a receptor is the progesterone receptor mediated by the steroid progesterone. At the progesterone receptor, progesterone acts as a natural agonist whereas it displays potent antimineralocorticoid properties both at the molecular and the systemic level. Besides classical effects on the uterus, antiepileptic, anxiolytic, hypnotic and anesthetic properties have been attributed to progesterone according to numerous studies.

Methods have been proposed for the use of mutant receptors, including mutant steroid receptors for gene therapy. For example, such methods are disclosed in WO 93/23431, WO 98/18925, WO 96/40911. WO 98/33903 discloses a genetic construct

comprising a steroid responsive element from a tissue specific gene, a coding sequence, and an SV40 enhancer.

Brief Description of the Invention

5 The object of the present invention is to overcome the disadvantages of the previous gene therapy delivery systems. The delivery system of the present invention is a composition of matter comprising a nucleic acid comprising at least one hormone responsive element (HRE), said hormone responsive element being coupled to a hormone-hormone receptor complex. A preferred embodiment of the composition of
10 matter of the invention is one where the hormone responsive element is a steroid responsive element (SRE), and the receptor is a steroid receptor. Most preferably, the hormone responsive element is a progesterone responsive element (PRE), and the receptor is a progesterone receptor.

15 The present invention provides a delivery system for gene therapy that should overcome the prior art disadvantages. The presence of the hormone responsive element together with the nucleic acid encoding a gene of interest stimulates and enhances gene expression and, more importantly, encourages the binding of a hormone-hormone receptor complex. One hormone responsive element is preferably present as a nucleic acid dimer sequence or nucleic acid multimer sequence. Even an inverse orientation of
20 the hormone responsive element will exert its proper function. The hormone-hormone receptor complex contains a hormone receptor that becomes activated after binding of its specific hormone. The hormone receptor in the activated state is able to recognize and bind to its specific hormone responsive element, which in the present invention is combined with a nucleic acid encoding the desired gene, e.g., a human blood-clotting
25 factor.

Another aspect of the present invention is a nucleic acid construct comprising at least one hormone responsive element (HRE), and in particular, comprising at least one hormone responsive element for regulating the expression of a gene encoding a human blood clotting factor. A preferred embodiment is one where the hormone responsive

element is a steroid responsive element (SRE). Most preferably, the hormone responsive element is a progesterone responsive element (PRE).

Another aspect of the present invention are vectors comprising the nucleic acid constructs of the present invention. Embodiments of the invention further include
5 transfected and transformed cells comprising these vectors and/or nucleic acids.

A further embodiment of the present invention is pharmaceutical compositions comprising a therapeutically effective dose of the nucleic acid constructs of the invention and a hormone. The hormone is preferably a steroid, and most preferably, progesterone. Another aspect of the invention is the use of the compositions of matter
10 and nucleic acid constructs as medicaments against genetic disorders or diseases, such as hemophilia, as well as the use of the compositions of matter and nucleic acid constructs for the manufacture of a medicament against genetic disorders and diseases, such as hemophilia.

The invention further includes a method of introducing into a cell a nucleic acid
15 construct encoding a gene of interest, e.g., a human blood-clotting factor, to express the gene of interest in the cell. This method incorporates a nucleic acid into the cell (via, for example, a vector) so that the cell expresses the gene encoded by the foreign nucleic acid. In this method, the nucleic acid encoding the gene, e.g., a human blood-clotting factor, is combined with a nucleic acid construct comprising at least one hormone
20 responsive element (HRE), preferably a progesterone responsive element. The presence of the hormone responsive element together with the nucleic acid encoding the gene, e.g., a human blood-clotting factor, stimulates and enhances gene expression and encourages the binding of a hormone-hormone receptor complex in the target cell.

Another aspect of the invention is a method of treating a blood clotting disorder
25 by administering a therapeutically effective amount of the composition of matter of the invention to an organism.

According to the method of the invention vaccination is another aspect of the embodiment. Introducing a nucleic acid construct or composition of matter of the invention comprising a gene for an antigen or containing viral sequence into a cell (DNA

or mRNA vaccines) using the method mentioned above may also provide a way to stimulate the cellular immune response.

Brief Description of the Drawings

5 Figure 1 is a diagram of the vector pTGFG1.

Figure 2 is a diagram of the vector pTGFG4.

Figure 3 is a diagram of the vector pTGFG5.

Figure 4 is a diagram of the vector pTGFG6.

Figure 5 is a diagram of the vector pTGFG7.

10 Figure 6 is a diagram of the vector pTGFG8.

Figure 7 is a diagram of the vector pTGFG9.

Figure 8 is a diagram of the vector pTGFG10.

Figure 9 is a diagram of the vector pTGFG11.

Figure 10 is a diagram of the vector pTGFG13.

15 Figure 11 is a diagram of the vector pTGFG14.

Figure 12 is a diagram of the vector pTGFG15.

Figure 13 is a diagram of the vector pTGFG16.

Figure 14 is a diagram of the vector pTGFG18.

Figure 15 is a diagram of the vector pTGFG19.

20 Figure 16 is a diagram of the vector pTGFG20.

Figure 17 is a diagram of the vector pTGFG21.

Figure 18 is a diagram of the vector pTGFG22.

Figure 19 is a diagram of the vector pTGFG23.

Figure 20 is a diagram of the vector pTGFG24.

Figure 21 is a diagram of the vector pTGFG25.

5 Figure 22 is a diagram of the vector pTGFG26.

Figure 23 is a diagram of the vector pTGFG27.

Figure 24 is a diagram of the vector pTGFG28.

Figure 25 is a diagram of the vector pTGFG29.

Figure 26 is a diagram of the vector pTGFG30.

10 Figure 27 is a diagram of the vector pTGFG31.

Figure 28 is a diagram of the vector pTGFG32.

Figure 29 is a diagram of the vector pTGFG33.

Figure 30 is a diagram of the vector pTGFG2.

Figure 31 is a diagram of the vector pTGFG34.

15 Figure 32 is a diagram of the vector pTGFG35.

Figure 33 is a diagram of the vector pTGFG36.

Figure 34 is a diagram of the vector pTGFG37.

Figure 35 is a diagram of the vector pTGFG38.

Figure 36 is a diagram of the vector pTGFG53.

20 Figure 37 is a diagram of the vector pTGFG64.

Figure 38 is a diagram of the vector pTGFG0.

Figure 39 is a diagram of the vector pTGFG66.

Figure 40 is a diagram of the vector pTGFG67.

Figure 41 is a diagram of the vector pTGFG68.

5 Figure 42 is a diagram of the vector pTGFG69.

Figure 43 is a diagram of the vector pTGFG82.

Figure 44 is a diagram of the vector pTGFG95.

Figure 45 is the DNA sequence of vector pTGFG36 (SEQ ID No. 1).

Figure 46 is the DNA sequence of vector pTGFG67 (SEQ ID No. 2).

10 Figure 47 shows a GFP concentration curve for cell homogenates after transfection with pTGFG5 and pTGFG20, respectively.

Figure 48 shows corresponding light (a and c) and fluorescent (b and d) micrographs of HeLa cells transfected with pTGFG5 (a and b) and pTGFG20 (c and d), respectively.

15

Detailed Description of the Invention

1. Definitions

"Nucleic acid" means DNA, cDNA, mRNA, tRNA. The nucleic acid may be
20 linear or circular, double-stranded or single-stranded.

"Nucleic acid construct" refers to a composite of nucleic acid elements in relation to one another. The nucleic acid elements of the construct may be incorporated into a

vector in such an orientation that a desired gene may be transcribed, and if desired, a desired protein may be expressed.

"Hormone responsive element" (HRE) refers to regions of nucleic acids, and in particular, DNA, which regulate transcription of genes in response to hormone activation. HREs are typically about 10-40 nucleotides in length, and more usually, about 13-20 nucleotides in length. As explained above, HREs become activated when a hormone binds to its corresponding intracellular receptor causing a conformational change, so that the receptor has increased affinity for the HRE and binds to it. The HRE, in turn, stimulates transcription. A "steroid-responsive element" (SRE) is an HRE that regulates transcription of genes in response to steroid activation. A "progesterone responsive element" (PRE) is an HRE/SRE that regulates transcription of genes in response to progesterone activation.

A "hormone receptor" refers to a receptor which binds to or is activated by a hormone. A "steroid receptor" refers to a receptor which binds to or is activated by a steroid hormone. A "progesterone receptor" is a receptor which binds to or is activated by the steroid hormone progesterone.

"Gene" refers to DNA involved in expressing a polypeptide, optionally including leader and trailer sequences and introns and exons.

"Vector" refers to any genetic construct, such as a plasmid, phage, cosmid, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. The term includes cloning and expression vehicles.

"Promoter" refers to a region of regulatory DNA sequences for the control of transcription of a gene to which RNA polymerase binds. The promoter forms an initiation complex with RNA polymerase to initiate and drive transcription activity. "Enhancers" may activate the complex or "silencers" may inhibit the complex. A "tissue-specific promoter" is a promoter found in the DNA of tissue for transcription of genes expressed in specific tissue.

"Therapeutically effective dose" of the products of the invention refers to a dose effective for treatment or prophylaxis, for example, a dose that yields effective treatment or reduction of the symptoms of hemophilia. It is also a dose that measurably activates expression of a target gene as determined by measurements of target protein levels, or a dose that is predictable to be effective for treatment or prophylaxis by extrapolating from *in vitro* or *in vivo* data. The determination of a therapeutically effective dose is within the purview of one skilled in the art.

"Encodes" or "encoding" refers to a nucleic acid sequence transcribed (in case of DNA) or translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences.

For the purposes of this application, "express", "expressing" or "expression" shall refer to transcription and translation of a gene encoding a protein.

2. Detailed Description and Examples

As stated above, an object of the present invention is to provide a new and improved delivery system for gene therapy. The invention thus provides in one embodiment a composition of matter comprising a nucleic acid comprising at least one hormone responsive element (HRE), said hormone responsive element being coupled to a hormone-hormone receptor complex. A preferred embodiment of the composition of matter of the invention is one where the hormone responsive element is a steroid responsive element (SRE), and the receptor is a steroid receptor. Most preferably, the hormone responsive element is a progesterone responsive element (PRE), and the receptor is a progesterone receptor.

Potential HREs for use in the present invention have been previously described. For example, GREs (Scheidereit, C., et al., *Nature*, Vol. 304, 749, 1983; von der Ahe, D., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 83, 2817, 1986), EREs or PREs (Chambon, P., et al., *Rec. Prog. Horm. Res.*, Vol., 40, 1, 1984; Klock, G., et al., *Nature*, Vol. 329, 734, 1987). As already stated above, the most preferred HRE for the invention is a PRE.

Specifically, the preferred PRE is described in Example I. The nucleic acid for use in the invention comprises at least one hormone responsive element. Preferred is a nucleic acid comprising one HRE, but it may comprise more than one HRE. For example, the nucleic acid may comprise three or five HREs. The most preferred embodiment is a
 5 nucleic acid comprising one PRE.

Potential hormone receptors for use in the present invention are, for example, estrogen receptors, mineralocorticoid receptors, glucocorticoid receptors, retinoic acid receptors, androgen, calcitriol, thyroid hormone or progesterone receptors and orphan receptors. Such receptors have been previously described. (Green, S., et al., *Nature*, Vol. 320, 134, 1986; Green, G. L., et al., *Science*, Vol. 231, 1150, 1986; Arriza, J. L., et al., *Science*, Vol. 237, 268, 1987; Hollenberg, S. M., et al., *Nature*, Vol. 318, 635, 1985; Petkovitch, M., et al., *Nature*, Vol. 330, 444, 1987; Giguere, V., et al., *Nature*, Vol. 330, 624, 1987; Tilley, W., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 86, 327, 1989; Baker, A. R., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 85, 3294, 1988; Weinberger, C., et al.,
 10 *Nature*, Vol. 324, 641, 1986; Sap, J., et al., *Nature*, Vol. 324, 635, 1086; Misrahi, M., et al., *Biochem. Biophys. Res. Commun.*, Vol. 143, 740, 1987; Kastner, P., et al., *Cell*, Vol. 83, 859, 1995). These receptors may be from human or other mammalian sources, although human is preferred. Nucleotide and/or amino acid sequences of human steroid receptors are available in the GenBank: mineralocorticoid receptor: M16801;
 15 glucocorticoid receptor α : M10901; glucocorticoid receptor α_2 : U01351; glucocorticoid receptor β : M11050; retinoic acid receptor α : AF088888 (exon 1), AF088889 (exon 2), AF088890 (exon 3), AF088891 (exon 4), AF088892 (exon 5 and 6), AF088893 (exon 7), AF088894 (exon 8), AF088895 (exon 9 and complete cDNA); retinoic acid receptor γ : M24857; androgen receptor: M27423 (exon 1), M27424 (exon 2), M27425 (exon 3),
 20 M27436 (exon 4), M27427 (exon 5), M27428 (exon 6), M27429 (exon 7), M27430 (exon 8); thyroid hormone receptor α_1 : M24748, thyroid hormone receptor α_2 : J03239; progesterone receptor: AF016381; somatotropin receptor: J00148; vitamin D receptor (calcitriol receptor): J03258.
 25

The skilled person will understand that expression of the receptor proteins can be
 30 achieved by standard methods, e.g. via PCR-cloning of the known cDNAs from cDNA libraries and overexpression of the corresponding proteins in suitable expression vectors,

such as, for example, the vectors of the present invention, in suitable host cells, e.g., COS cells. Accordingly, subsequent purification of the cytosolic fraction can be achieved by routine methods such as affinity chromatography purification. For this purpose, various suitable antibodies against the desired receptor are commercially available. For example, polyclonal antibodies against the mouse progesterone receptor that have a sufficiently high cross-reactivity for the human protein are available from Dianova (Hamburg, Germany). Likewise, further purification can be achieved by standard methods, e.g., chromatographical methods such as ion-exchange chromatography and/or FPLC.

The most preferred receptor is the progesterone receptor. Preferably, the receptor is a human progesterone receptor. Such a human progesterone receptor (from T47D human breast cancer cells) is disclosed in US Patent No. 4,742,000, and cells expressing this receptor have been deposited (ATCC deposit number HTB, 133). As already described above, it would be routine to purify such a receptor from the cytosol using receptor specific antibodies. In addition, US Patent No. 4,742,000 discloses a method for purification of the human progesterone receptor using a specific steroid affinity resin (cf. Grandics et al., Endocrinology, Vol. 110, 1088, 1982). Briefly, the cytosolic fraction of the T47D cells is passed over Sterogel, a commercial preparation of deoxycorticosterone coupled to Sepharose 2B that selectively binds the progesterone receptor. After washing with loading buffer, the bound receptor is eluted with a buffer containing progesterone. The eluted steroid-receptor complex is then chromatographed on DEAE-Biogel and eluted stepwise with a buffer containing 0.2M NaCl. Subsequently, the bound progesterone can be readily exchanged. As described above, further purification can be achieved by routine methods well-known to the skilled person.

Mutated versions of these receptors and derivatives thereof, that still retain the function of the receptors to bind a ligand and thereby become activated and bind DNA and regulate transcription, may also be employed in the invention. Such derivative may be a chemical derivative, variant, chimera, hybrid, analog, or fusion.

The hormone in the composition of matter may include synthetic and natural hormones, such as estrogen, testosterone, glucocorticoid, androgen, thyroid hormone, and progesterone or derivatives thereof. These are widely available. Progesterone is most preferred. For example, natural micronized progesterone has been marketed in France under the trademark of UTROGESTAN® since 1980. Its properties are similar to the endogenous progesterone, in particular, it has antiestrogen, gestagen, slightly antiandrogen and antimineralocorticoid properties.

Micronized progesterone has advantages that make it a suitable carrier for genes or nucleic acid constructs to target cells. Specifically, the synergistic effect of the double process of micronization and suspension in long-chain fatty acids results in increasing progesterone absorption. It has been demonstrated that after oral administration of 100 mg of UTROGESTAN®, peak plasma progesterone levels were obtained after 1-4 hours in most cases (Padwick, M. L., et al., *Fertil. Steril.*, Vol. 46, 402, 1986). Later on, the levels declined substantially, although they were still elevated at 12 hours. Even at 84 hours the levels were slightly higher than baseline. A U.S. kinetic study confirmed earlier work demonstrating the bioavailability of oral micronized progesterone. They showed a peak effect at 2 hours followed by rapid decrease in plasma progesterone level (Simon, J. A., et al., *Fertil., Steril.*, Vol., 60, 26, 1993).

A further advantage of using progesterone as a carrier is the low level of disadvantageous side effects. Orally administered progesterone adversely affects neither plasma lipids (Jensen, J. et al., *Am. J. Obstet. Gynecol.*, Vol. 156, 66, 1987) nor carbohydrate metabolism (Mosnier-Pudar, H. et al., *Arch. Mal. Coeur*, Vol 84, 1111, 1991). Further, progesterone does not affect liver enzymes (ASAT, ALAT, AFOS), sex-hormone binding-globulin (SHBG) synthesis or HDL-cholesterol levels at daily doses of 200 mg and 300 mg. Although the plasma levels of deoxycorticosterone may increase substantially during UTROGESTAN® treatment, there are strong indications that the mineralocorticoid effects of this progesterone metabolite are completely counteracted by the anti-mineralocorticoid effects of progesterone itself. This is apparent from a comparative study (Corvol, P., et al., In: *Progesterone and progestins*. Raven Press, New York, 179, 1983) in which oral UTROGESTAN® was capable of antagonizing the mineralocorticoid effects of 9-alpha-fluorohydrocortisone.

The skilled artisan will appreciate that the composition may contain other components capable of assisting in introducing the nucleic acid into a cell for the purpose of gene therapy. Specifically, the composition may further contain β -cyclodextrine, glycerin, lecithin or corn oil. For example, the composition of hormone-hormone receptor nucleic acid complex of the invention may be provided orally to humans or animals as a gelatin capsule. Progesterone therein could be present in a concentration of 200 -300 mg dissolved in a 35 % or 40 % β -cyclodextrin solution or in cornoil or glycerol with peanut oil together with lecithin.

Another aspect of the present invention is a nucleic acid construct comprising at least one hormone responsive element (HRE), and in particular, comprising at least one hormone responsive element for regulating the expression of a gene, such as, for example, a gene encoding a human blood clotting factor. A preferred embodiment is one where the hormone responsive element is a steroid responsive element (SRE). Most preferably, the hormone responsive element is a progesterone responsive element (PRE). This nucleic acid construct may be used as the nucleic acid in the composition of matter of the first aspect of the invention.

Aside from the HREs, SREs, or PREs already disclosed above, the nucleic acid in the present invention may further contain promoter, enhancer, or silencer sequences. The promoter may be ubiquitous or tissue-specific. Of the ubiquitous promoters, the CMV promoter is most preferred. However, a tissue-specific promoter is preferred over a ubiquitous promoter. For example, the tissue-specific promoters envisioned for the instant invention include α_1 -antitrypsin. The nucleic acid construct may further comprise additional sequences such as the ampicillin resistance gene. Other reporter sequences known to the skilled artisan may also be included, such as, for example, the green fluorescent protein (GFP), luciferase, β -galactosidase or chloramphenicol-transferase (CAT). As an enhancer sequence, the SV40 intron and SV40 Poly A are most preferred. A preferred nucleic acid construct contains sequentially from the 5' to the 3' end: a PRE, a CMV promoter, a gene of interest, SV40 Intron and SV40 poly A enhancer sequence, and an ampicillin resistant gene.

The gene of interest may be chosen from those encoding proteins lacking in a variety of genetic disorders or involved in conditions related to inappropriate responses to hormones, for example, hormone-dependent cancers such as breast, ovarian, and endometrial cancers and prostate cancer. The gene of interest may also be used to replace a defective gene resulting in such genetic disorders as hemophilia, von Willebrand disease, and cystic fibrosis. The gene of interest includes mutations of such gene or a gene encoding a fusion product. The nucleic acid construct of the present invention may comprise more than one gene of interest.

In particular, the gene of interest may replace genes for a blood clotting factor, and preferably a human blood-clotting factor. The genes encoding Factor VIII and Factor IX, involved in Hemophilia A and B, respectively, are good candidates for the invention. Other candidates include the gene encoding von Willebrand factor, Factor IV, Factor X, or protein C.

Other useful genes include, but are not limited to, hormone genes such as the genes encoding for insulin, parathyroid hormone, luteinizing hormone releasing factor (LHRH), alpha and beta seminal inhibins and human growth hormone; hormone receptor genes such as the glucocorticoid receptor, the estrogen receptor, the progesterone receptor, the retinoic acid receptor; growth factors such as vascular endothelial growth factor (VEGF), nerve growth factor, epidermal growth factor; enzyme genes; genes encoding cytokines or lymphokines such as interferons, granulocytic macrophage colony stimulating factor (GM-CSF), colony stimulating factor-1 (CSF-1), tumor necrosis factor (TNF), and erythropoietin (EPO); genes encoding inhibitor substances such as alpha₁-antitrypsin, and genes encoding substances that function as drugs, e. g., genes encoding the diphtheria and cholera toxins, ricin or cobra venom factor. Also, antisense sequences may be administered as genetic material.

Another aspect of the present invention is vectors comprising the nucleic acid constructs of the present invention. These vectors may be used in the composition matter of the present invention. Preferably, however, the nucleic acid sequence for use in the invention is linear rather than circular. The vectors may be capable of expressing

Embodiments of the invention further include transfected and transformed cells comprising these vectors and/or nucleic acid constructs. Within the scope of this invention, a transfected cell is one in which foreign DNA has been incorporated. Methods of transfection may include microinjection, CaPO_4 precipitation, electroporation, liposome fusion, or by gene gun. Most preferably, transfection is achieved by electroporation.

A further embodiment of the present invention is pharmaceutical compositions comprising a therapeutically effective dose of the nucleic acid constructs of the invention and a hormone. The hormone is preferably a steroid, and most preferably, progesterone, as described above. The dose is dependent on the condition to be treated, the characteristics of the patient, and the result sought to be achieved. Determining dosage is within the realm of the skilled artisan.

The invention further includes a method of introducing into a cell a nucleic acid construct encoding a gene of interest, e.g., a human blood-clotting factor, to express the blood-clotting factor in the cell. In this method, the nucleic acid encoding a human blood-clotting factor is combined with a nucleic acid construct comprising at least one hormone responsive element (HRE), preferably a progesterone responsive element.

The mixture of nucleic acid bound to the hormone-hormone receptor complex together with an excess of hormone, preferably progesterone, will be used to introduce the nucleic acid into a cell by various methods known to the skilled artisan and outlined above. The cell-uptake will be stimulated by the interaction of the hormone at the cell
5 membrane level. The hormone or steroid interacts with the lipid bilayer of the cell membrane not only through membrane perturbation but also through activation of certain hormone- or steroid-sensitive membrane receptors. This has been demonstrated for progesterone and other steroids. Last but not least, it is known that hormones are able to cross the cell membrane by diffusion. In the present invention, the nucleic acid bound
10 to the hormone-hormone receptor complex should be transported through the membrane during the process of diffusion.

Another aspect of the invention is a method of treating a blood clotting disorder by administering a therapeutically effective amount of the composition of matter of the invention to an organism. This method involves the administration and dosage
15 considerations already discussed.

Experiments have been performed to illustrate the technical aspects of the present invention. These experiments are described in Examples II-IV.

The following are examples of the present invention. The skilled artisan will be readily recognize that the invention is not limited to these examples.

20

Example I: Construction of Vectors

Production of the vector pTGFG1

The vector pUC19 (MBI Fermentas) was digested with XbaI, treated with
25 Klenow-Enzyme and religated. This XbaI deleted vector was then digested with EcoRI, treated with Klenow-Enzyme and religated in order to delete the EcoRI site. For insertion of a XbaI site in the SacI site of this vector it was digested with SacI, treated

with T4-Polymerase, dephosphorylated with Alkaline Phosphatase and ligated with the XbaI-linker CTCTAGAG (Biolabs #1032). Another XbaI-site was inserted by digesting the newly produced vector with HindIII, treating it with Klenow, dephosphorylating it with Alkaline Phosphatase and ligating it with the XbaI-linker CTCTAGAG (Biolabs #1032). This vector was named pUC19/X.

In order to destroy the XbaI-site present in the vector pHGF-S65T (Clontech) this vector was digested with XbaI, treated with Klenow-Enzyme and religated resulting in the vector pGFP/0. A 2.3 kb fragment containing the GFP-Gene was isolated after digesting pGFP/0 with MluI, treating it with Klenow-Enzyme and digesting it with BamHI. This fragment was inserted into the multiple cloning site of the vector pUC19/X which was digested with Sall, treated with Klenow-Enzyme and digested with BamHI. The resulting vector was named pTGFG1 (Figure 1).

Production of the insert PRE(ds)

The Oligonucleotides (Metabion) PRE-S (5'-GGG GTA CCA GCT TCG TAG CTA GAA CAT CAT GTT CTG GGA TAT CAG CTT CGT AGC TAG AAC ATC ATG TTC TGG TAC CCC-3') (SEQ ID No. 3) and PRE-AS (5'-GGG GTA CCA GAA CAT GAT GTT CTA GCT ACG AAG CTG ATA TCC CAG AAC ATG ATG TTC TAG CTA CGA AGC TGG TAC CCC-3') (SEQ. ID No. 4) were hybridized and phosphorylated by kinase reaction, resulting in the inserts PRE(ds).

Production of the vector pTGFG4

The vector pTGFG1 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG4 (Figure 2).

Production of the vector pTGFG5

The vector pTGFG1 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG5 (Figure 3).

Production of the vector pTGFG6

The vector pTGFG1 was digested with *EheI* and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG6 (Figure 4).

5 Production of the vector pTGFG7

The vector pTGFG1 was digested with *KpnI*, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG7 (Figure 5).

Production of the vector pTGFG8

10 The vector pTGFG7 was digested with *EheI* and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG8 (Figure 6).

Production of the vector pTGFG9

15 The vector pTGFG7 was digested with *SapI*, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG9 (Figure 7).

Production of the vector pTGFG10

20 The vector pTGFG7 was digested with *EcoO109I*, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG10 (Figure 8).

Production of the vector pTGFG11

The vector pTGFG7 was digested with *PstI*, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG11. (Figure 9).



Production of the vector pTGFG13

The vector pTGFG6 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG13 (Figure 10).

5 Production of the vector pTGFG14

The vector pTGFG6 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG14 (Figure 11).

Production of the vector pTGFG15

10 The vector pTGFG5 was digested with SapI, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG15 (Figure12).

Production of the vector pTGFG16

15 The vector pTGFG5 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG16 (Figure 13).

Production of the vector pTGFG18

20 The vector pTGFG9 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG18 (Figure 14).

Production of the vector pTGFG19

The vector pTGFG10 was digested with EheI and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG19 (Figure 15).

Production of the vector pTGFG20

The vector pTGFG11 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG20 (Figure 16).

5 Production of the vector pTGFG21

The vector pTGFG15 was digested with EheI and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG21 (Figure 17).

Production of the vector pTGFG22

10 The vector pTGFG14 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG22 (Figure 18).

Production of the vector pTGFG23

15 The vector pTGFG11 was digested with EheI and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG23 (Figure 19).

Production of the vector pTGFG24

20 The vector pTGFG9 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG24 (Figure 20).

Production of the vector pTGFG25

The vector pTGFG14 was digested with SapI, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG25 (Figure 21).

Production of the vector pTGFG26

The vector pTGFG16 was digested with SapI, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG26 (Figure 22).

5 Production of the vector pTGFG27

The vector pTGFG9 was digested with EheI and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG27 (Figure 23).

Production of the vector pTGFG28

- 10 The vector pTGFG18 was digested with EheI and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG28 (Figure 24).

Production of the vector pTGFG29

- 15 The vector pTGFG25 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG29 (Figure 25).

Production of the vector pTGFG30

- 20 The vector pTGFG27 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG30 (Figure 26).

Production of the vector pTGFG31

The vector pTGFG24 was digested with EcoO109I, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG31 (Figure 27).

Production of the vector pTGFG32

The vector pTGFG19 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG32 (Figure 28).

5 Production of the vector pTGFG33

The vector pTGFG28 was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG33 (Figure 29).

Production of the vector pTGFG2

10 The vector pUC19 (MBI Fermentas) was digested with SalI, treated with Klenow-Enzyme and dephosphorylated with Alkaline Phosphatase. It was ligated to the NotI-linker GCGGCCGC (Biolabs # 1045), resulting in the vector pUC19/N.

A 1.4 kb fragment containing the open reading frame of the human clotting factor IX, isolated from a human cDNA library, was inserted into the PstI-site of the vector
 15 pUC19/N which was digested with PstI, treated with T4-Polymerase and dephosphorylated with Alkaline Phosphatase. From the resulting vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG1. The resulting
 20 vector was named pTGFG2 (Figure 30).

Production of the vector pTGFG34

From the vector pUC19/N-FIX, a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with HindIII and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-
 25 digested vector pTGFG3 resulting in the vector pTGFG34 (Figure 31).

Production of the vector pTGFG35

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with HindIII and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG4 resulting in the vector pTGFG35 (Figure 32).

Production of the vector pTGFG36

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG5 resulting in the vector pTGFG36 (Figure 33). This vector is a preferred one for delivery of Factor IX into the cell, and its DNA sequence is provided in Figure 46.

Production of the vector pTGFG37

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with HindIII and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG6 resulting in the vector pTGFG37 (Figure 35).

Production of the vector pTGFG38

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG7 resulting in the vector pTGFG38 (Figure 35).

Production of the vector pTGFG53

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG20 resulting in the vector pTGFG53 (Figure 36).

Production of the vector pTGFG64

From the vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG33 resulting in the vector pTGFG64 (Figure 37).

Production of the insert ALLG(ds)

The Oligonucleotides (Metabion) ALLG1/1 (5'-AGC TTG ACC TCG AGC AAG C-3') (SEQ. ID NO: 6) and ALLG2 (5'-GGC CGC TTG CTC GAG GTC A-3') (SEQ. ID NO: 7) were hybridized and phosphorylated by kinase reaction, resulting in the inserts ALLG(ds). The insert ALLG (ds) was constructed to introduce into the vector of choice a sequence with a multiple cloning site for the possible introduction of other transgenes.

Production of the vector pTGFG0

The vector pTGFG1 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG0 (Figure 38).

Production of the vector pTGFG66

The vector pTGFG4 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG66 (Figure 39).

Production of the vector pTGFG67

The vector pTGFG5 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG67 (Figure 40). This vector is preferred for the delivery of any gene inserted into the cloning site, and its DNA sequence is provided in Figure 47.

Production of the vector pTGFG68

The vector pTGFG6 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG68 (Figure 41).

5 Production of the vector pTGFG69

The vector pTGFG7 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG69 (Figure 42).

Production of the vector pTGFG82

10 The vector pTGFG20 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG82 (Figure 43).

Production of the vector pTGFG95

15 The vector pTGFG33 was double-digested with HindIII and NotI. The 4.3 kb fragment was ligated with the insert ALLG(ds), resulting in the vector pTGFG95. (Figure 44)

Example II: Isolation of Human Factor IX cDNA

Factor IX cDNA was amplified from human liver cDNA (Clontech) using two
20 primers overlapping the start and termination codon of the factor IX open reading frame resulting in a 1387 bp fragment containing the entire open reading frame. Restriction sites for EcoRI (upstream) and BamHI (downstream) were included at the end of each primer to facilitate cloning. Amplification was performed with Pwo Polymerase (Boehringer Mannheim) in 50 µl reaction volume [10 mM Tris HCl pH 8.85, 25 mM
25 KCl, 5 mM (NH₄)₂SO₄, 2 mM MgSO₄] with 30 incubation cycles 96° C 1 min., 60° C 1 min., 72° C 2 min., followed by a final extension step at 72° C 10 min.

Reaction products were ligated into the EcoRI- and BamHI-sites of pUC19 and transformed into *E. coli* DH5-a. Positive clones were selected. Sequences were confirmed by cycle sequencing (Amersham) from both ends with labeled primers (IR-700) and automated analysis on the LiCor sequencing system (MWG, Biotech).

5 The following primers were used :

gGAATTCcgcaaagggttATGCAGCGCGTGAACATGATCATGGC(upstream)(SEQ. ID NO:8)

cgcGGATCCATTAAGTGAGCTTTGTTTTTTCCTTAATCC (downstream)(SEQ. ID NO:9)

10

Example III: Expression and Quantification of the Marker Protein GFP ("Green Fluorescent Protein")

15 HeLa cells were transfected by electroporation with plasmids pTGFG5 or pTGFG20. Transfected cells were harvested, and the cell pellets were homogenized and lysated in a buffer containing phosphate-buffered saline (pH 7.5) and 10 mM PMSF. The concentration of green fluorescent protein (GFP) in the cell homogenate was determined by competitive ELISA.

20 For this purpose, GFP was coated in a defined concentration on microtiter plates. Then, GFP samples were added in presence of anti-GFP antibody. After several washing steps, a labeled secondary antibody was added in order to trace the first antibody. The colorimetric reaction was measured photometrically (extinction). Generally, the more GFP was added, the less antibody was left to bind the coated GFP. Thus, reduction of extinction corresponded to higher GFP concentration in the sample.

25 A concentration curve of GFP was determined by linear regression (Figure 47) using bovine serum albumin (BSA) as a reference. A mean value of 2.4 µg GFP/ml for pTGFG5 (1 PRE) and 5.2 µg GFP/ml for pTGFG20 (3PREs) was found.

Figures 48 a-d show micrographs of HeLa cell cultures transfected with pTGFG5 (Fig. 48 a and b) and pTGFG20 (Fig. 48 c and d), respectively. Figures 46 a and c represent light microscopic views as controls, and Fig. 48 b and d show the corresponding cell patches in the fluorescent mode. Routinely, more than 50% of the cells expressed GFP, indicating very efficient expression, the presence of only one PRE showing more efficient expression.

Example IV: Human Factor IX Quantification by ELISA Assay

HeLa cells were transfected either by electroporation or using liposome reagent DOTAP (Boehringer Mannheim) with plasmids pTGFG36, pTGFG53 and pTGFG64. These plasmids contain the cDNA of human clotting factor IX. Recombinant human factor IX was secreted into the supernatant of the cell culture and quantified using a sandwich ELISA method.

0.11 M sodium citrate and 10 mM PMSF were added in order to prevent degradation of human factor IX. The enzyme-immunological in vitro assay "Asserachrom IX:AG" from Boehringer-Mannheim was used in order to determine the concentration of expressed human clotting factor IX. The factor IX-standard from Octapharma AG was used as a standard in aqueous solutions of 28 IU/ml.

In six different transfection experiments, in which HeLa cells with plasmids containing human factor IX-cDNA (pTGFG36, 53 and 64) were transfected using either electroporation or lipid-transfection reagent (DOTAP, Boehringer Mannheim), a concentration range of 3-25 ng/ml human clotting factor IX was reached.

Claims:

1. A composition of matter comprising a nucleic acid comprising a hormone responsive element (HRE), said hormone responsive element being coupled to a hormone-hormone receptor complex.
- 5 2. The composition of matter of claim 1, wherein the nucleic acid further comprises a gene encoding a blood clotting factor.
3. The composition of matter of claim 2, wherein the human blood clotting factor is selected from the group consisting of Factor VIII, Factor IX, and von Willebrand Factor (vWF).
- 10 4. The composition of matter of claim 2, wherein the hormone responsive element is a steroid responsive element (SRE).
5. The composition of matter of claim 4, wherein the steroid responsive element (SRE) is a progesterone responsive element (PRE).
- 15 6. The composition of matter of claim 1, wherein the complex is a steroid-steroid receptor complex.
7. The composition of matter of claim 6, wherein the receptor is a progesterone receptor and the steroid is progesterone.
8. The composition of matter of claim 3, where the human blood clotting factor is Factor IX.
- 20 9. The composition of matter of claim 7, where the human blood clotting factor is Factor IX.
10. A nucleic acid construct comprising at least one hormone responsive element (HRE) for regulating the expression of a gene encoding a human blood clotting factor.

11. The nucleic acid construct of claim 10, wherein the hormone responsive element is a steroid responsive element (SRE).
12. The nucleic acid construct of claim 11, wherein the steroid responsive element is a progesterone responsive element (PRE).
- 5 13. The nucleic acid construct of claim 10, wherein the human blood clotting factor is selected from the group consisting of Factor VIII, Factor IX, and von Willebrand Factor (vWF).
14. The nucleic acid construct of claim 10, which further comprises a tissue-specific promoter.
- 10 15. The nucleic acid construct of claim 10, wherein the hormone responsive element (HRE) is a progesterone responsive element (PRE) and the blood clotting factor is Factor IX.
16. The nucleic acid construct of claim 15, which further comprises a tissue-specific promoter.
- 15 17. A vector comprising the nucleic acid construct of claim 10.
18. A vector comprising the nucleic acid construct of claim 11.
19. A vector comprising the nucleic acid construct of claim 12.
20. A vector comprising the nucleic acid construct of claim 13.
21. A vector comprising the nucleic acid construct of claim 14.
- 20 22. A vector comprising the nucleic acid construct of claim 15.
23. A vector comprising the nucleic acid construct of claim 16.
24. A cell transfected with the nucleic acid construct of claim 10.
25. A cell transformed by the nucleic acid construct of claim 10.

26. A pharmaceutical composition comprising a therapeutically effective dose of the nucleic acid construct of claim 10 and a hormone.
27. A pharmaceutical composition comprising a therapeutically effective dose of the nucleic acid construct of claim 15 and progesterone.
- 5 28. A pharmaceutical composition comprising a therapeutically effective dose of the composition of matter of claim 1.
29. A pharmaceutical composition comprising a therapeutically effective dose of the composition of matter of claim 9.
- 10 30. A method of delivering into a cell a nucleic acid encoding a gene to be expressed in the cell comprising providing the composition of matter of claim 1 to an organism so that the hormone in the composition crosses the cell membrane by diffusion and transports the nucleic acid that is coupled to the hormone-hormone receptor complex across the membrane and into the cell.
- 15 31. The method of claim 30, wherein a nucleic acid encoding human Factor IX is delivered into the cell.
32. A method of treating a blood clotting disorder comprising administering a therapeutically effective amount of the composition of matter of claim 1 to an organism.
- 20 33. A method of treating hemophilia B, comprising administering a therapeutically effective amount of the composition of matter of claim 9 to an organism.

Figure 1




Figure 2




Figure 3






Figure 4



19-02-99

4


Figure 5



19.02.99

40

Figure 6



19.02.2009

4

Figure 7




Figure 8





Figure 9



19.02.99

46

Figure 10




Figure 11




Figure 12




Figure 13




Figure 14





Figure 15



1999

Figure 16





Figure 17



199299

5

Figure 18





Figure 19



1999

5

Figure 20






Figure 21



19.02.99


54

Figure 22



19.02.99


Figure 23



19.02.99

6


Figure 24



19.02.99

6,

Figure 25



1999

6

Figure 26




Figure 27




Figure 28




Figure 29




Figure 30




Figure 31




Figure 32




Figure 33




Figure 34




Figure 35




Figure 36




Figure 37





Figure 38



19.02.99

7

Figure 39





Figure 40



19.02.99

Figure 41




Figure 42






Figure 43



19.09.99

Figure 44



1 seq pTGFG36	CGCGTTGACA TTGATTATTG ACTAGTTATT AATAGTAATC AATTACGGGG
1 seq pTGFG36	TCATTAGTTC ATAGCCCATATA TATGGAGTTC CGCGTTACAT AACTTACGGT
1 seq pTGFG36	AAATGGCCCG CCTGGCTGAC CGCCCAACGA CCCCCGCCCA TTGACGTCAA
1 seq pTGFG36	TAATGACGTA TGTTCACATA GTAACGCCAA TAGGGACTTT CCATTGACGT
1 seq pTGFG36	CAATGGGTGG AGTATTACG GTAAACTGCC CACTTGGCAG TACATCAAGT
1 seq pTGFG36	GTATCATATG CCAAGTACGC CCCCTATTGA CGTCAATGAC GGTAATGGC
1 seq pTGFG36	CCGCCTGGCA TTATGCCAG TACATGACCT TATGGACTT TCCTACTTGG
1 seq pTGFG36	CAGTACATCT ACGTATTAGT CATCGCTATT ACCATGGTGA TGCGGTTTIG
1 seq pTGFG36	GCAGTACATC AATGGGCGTG GATAGCGGTT TGA CTCACGG GGATTCCAA
1 seq pTGFG36	GTCTCCACCC CATTGACGTC AATGGGAGTT TGT TTTGGCA CAAAATCAA
1 seq pTGFG36	CGGGACTTTC CAAAATGTCG TAACAACTCC GCCCATTGA CGCAAATGGG
1 seq pTGFG36	CGGTAGGCGT GTACGGTGGG AGGTCTATAT AAGCAGAGCT CTCTGGCTAA
1 seq pTGFG36	CTAGAGAACC CACTGCTTAC TGGCTTATCG AAATTAATAC GACTCACTAT
1 seq pTGFG36	AGGGAGACCC AAGCTTGCAT GCCAATTCCG CAAAGGTTAT GCAGCGCGTG
1 seq pTGFG36	AACATGATCA TGGCAGAATC ACCAGGCCTC ATCACCATCT GCCTTTTAGG
1 seq pTGFG36	ATATCTACTC AGTGCTGAAT GTACAGTTTT TCTTGATCAT GAAAACGCCA
1 seq pTGFG36	ACAAAATTCT GAATCGGCCA AAGAGGTATA ATTCAGGTAA ATTGGAAGAG
1 seq pTGFG36	TTTGTTCAAG GGAACCTTGA GAGAGAATGT ATGGAAGAAA AGTGTAGTTT
1 seq pTGFG36	TGAAGAAGCA CGAGAAGTTT TTGAAAACAC TGAAAGAACA ACTGAATTTT

Sequenz pTGFG36 13.12.1998 20:16 Uhr

	960	970	980	990	1000
1 seq pTGFG36	GGAAGCAGTA	TGTTGATGGA	GATCAGTGTG	AGTCCAATCC	ATGTTTAAAT
	1010	1020	1030	1040	1050
1 seq pTGFG36	GGCGGCAGTT	GCAAGGATGA	CATTAATTC	TATGAATGTT	GGTGTCCTT
	1060	1070	1080	1090	1100
1 seq pTGFG36	TGGATTGAA	GGAAAGAACT	GTGAATTAGA	TGTAACATGT	AACATTAAGA
	1110	1120	1130	1140	1150
1 seq pTGFG36	ATGGCAGATG	CGAGCAGTTT	TGTAAAATA	GTGCTGATAA	CAAGGTGGTT
	1160	1170	1180	1190	1200
1 seq pTGFG36	TGCTCCTGTA	CTGAGGGATA	TCGACTTGCA	GAAAACCAGA	AGTCCTGTGA
	1210	1220	1230	1240	1250
1 seq pTGFG36	ACCAGCAGTG	CCATTTCAT	GTGGAAGAGT	TTCTGTTTCA	CAAACCTCTA
	1260	1270	1280	1290	1300
1 seq pTGFG36	AGCTCACCCG	TGCTGAGACT	GTTTTTCCTG	ATGTGGAATA	TGTAAATTCT
	1310	1320	1330	1340	1350
1 seq pTGFG36	ACTGAAGCTG	AAACCATTTT	GGATAACATC	ACTCAAAGCA	CCCAATCATT
	1360	1370	1380	1390	1400
1 seq pTGFG36	TAATGACTTC	ACTCGGGTTG	TTGGTGGAGA	AGATGCCAAA	CCAGGTCAAT
	1410	1420	1430	1440	1450
1 seq pTGFG36	TCCCTGGCA	GGTTGTTTTG	AATGGTAAAG	TTGATGCATT	CTGTGGAGGC
	1460	1470	1480	1490	1500
1 seq pTGFG36	TCTATCGTTA	ATGAAAATG	GATTGTAAC	GCTGCCCACT	GTGTTGAAAC
	1510	1520	1530	1540	1550
1 seq pTGFG36	TGGTGTTAAA	ATTACAGTTG	TCGCAGGTGA	ACATAATATT	GAGGAGACAG
	1560	1570	1580	1590	1600
1 seq pTGFG36	AACATACAGA	GCAAAGCGA	AATGTGATT	GAATTATTCC	TCACCACAAC
	1610	1620	1630	1640	1650
1 seq pTGFG36	TACAATGCAG	CTATTAATAA	GTACAACCAT	GACATTGCCC	TTCTGGAAC
	1660	1670	1680	1690	1700
1 seq pTGFG36	GGACGAACCC	TTAGTGCTAA	ACAGCTACGT	TACACCTATT	TGCATTGCTG
	1710	1720	1730	1740	1750
1 seq pTGFG36	ACAAGGAATA	CACGAACATC	TTCTCAAAT	TTGGATCTGG	CTATGTAAGT
	1760	1770	1780	1790	1800
1 seq pTGFG36	GGCTGGGGAA	GAGTCTTCCA	CAAAGGGAGA	TCAGCTTTAG	TTCTTCAGTA
	1810	1820	1830	1840	1850
1 seq pTGFG36	CCTTAGAGTT	CCACTTGTTG	ACCGAGCCAC	ATGTCTTCGA	TCTACAAAGT
	1860	1870	1880	1890	1900
1 seq pTGFG36	TCACCATCTA	TAACAACATG	TTCTGTGCTG	GCTTCCATGA	AGGAGGTAGA

Sequenz pTGFG36 13.12.1998 20:16 Uhr

10000

	1910	1920	1930	1940	1950
seq pTGFG36	GATTCATGTC	AAGGAGATAG	TGGGGGACCC	CATGTTACTG	AAGTGAAGG
	1960	1970	1980	1990	2000
seq pTGFG36	GACCAGTTTC	TTAACTGGAA	TTATTAGCTG	GGGTGAAGAG	TGTGCAATGA
	2010	2020	2030	2040	2050
seq pTGFG36	AAGGCAAATA	TGGAATATAT	ACCAAGGTAT	CCCGGTATGT	CAACTGGATT
	2060	2070	2080	2090	2100
seq pTGFG36	AAGGAAAAAA	CAAAGCTCAC	TTAATGGGAT	CGGTCGAGCG	GCCGCGACTC
	2110	2120	2130	2140	2150
seq pTGFG36	TACTAGAGGA	TCTTTGTGAA	GGAACCTTAC	TTCTGTGGTG	TGACATAATT
	2160	2170	2180	2190	2200
seq pTGFG36	GGACAAACTA	CCTACAGAGA	TTTAAAGCTC	TAAGGTAAAT	ATAAAATTTT
	2210	2220	2230	2240	2250
seq pTGFG36	TAAGTGTATA	ATGTGTTAAA	CTACTGATTC	TAATTGTTTG	TGTATTTTAG
	2260	2270	2280	2290	2300
seq pTGFG36	ATTCCAACCT	ATGGAAGTGA	TGAATGGGAG	CAGTGGTGGA	ATGCCTTTAA
	2310	2320	2330	2340	2350
seq pTGFG36	TGAGGAAAAC	CTGTTTGTCT	CAGAAGAAAT	GCCATCTAGT	GATGATGAGG
	2360	2370	2380	2390	2400
seq pTGFG36	CTACTGCTGA	CTCTCAACAT	TCTACTCCTC	CAAAAAAGAA	GAGAAAGGTA
	2410	2420	2430	2440	2450
seq pTGFG36	GAAGACCCCA	AGGACTTTCC	TTCAGAATIG	CTAAGTTTTT	TGAGTCATGC
	2460	2470	2480	2490	2500
seq pTGFG36	TGTGTTTAGT	AATAGAACTC	TTGCTTGCTT	TGCTATTTAC	ACCACAAAGG
	2510	2520	2530	2540	2550
seq pTGFG36	AAAAAGCTGC	ACTGCTATAC	AAGAAAATTA	TGGAAAAATA	TTCTGTAACC
	2560	2570	2580	2590	2600
seq pTGFG36	TTTATAAGTA	GGCATAACAG	TTATAATCAT	AACATACTGT	TTTTTCTTAC
	2610	2620	2630	2640	2650
seq pTGFG36	TCCACACAGG	CATAGAGTGT	CTGCTATTAA	TAACATGCT	CAAAAATTGT
	2660	2670	2680	2690	2700
seq pTGFG36	GTACCTTTAG	CTTTTAAATT	TGTAAAGGGG	TTAATAAGGA	ATATTTGATG
	2710	2720	2730	2740	2750
seq pTGFG36	TATAGTGCCT	TGACTAGAGA	TCATAATCAG	CCATACCACA	TTTGTAGAGG
	2760	2770	2780	2790	2800
seq pTGFG36	TTTTACTTGC	TTTAAAAAAC	CTCCACACCC	TCCCCCTGAA	CCTGAAACAT
	2810	2820	2830	2840	2850
seq pTGFG36	AAAATGAATG	CAATTGTTGT	TGTTAACTTG	TTTATTGCAG	CTTATAATGG

Sequenz pTGFG36 13.12.1998 20:16 Uhr

	2860	2870	2880	2890	2900
1 seq pTGFG36	TTACAAATAA	AGCAATAGCA	TCACAAATTT	CACAAATAAA	GCATTTTTTT
	2910	2920	2930	2940	2950
1 seq pTGFG36	CACTGCATTC	TAGTTGTGGT	TGTCCAAAC	TCATCAATGT	ATCTTATCAT
	2960	2970	2980	2990	3000
1 seq pTGFG36	GTCTGGATCC	CCGGGTACCC	TCTAGAGCGA	ATTAATTCAC	TGGCCGTCGT
	3010	3020	3030	3040	3050
1 seq pTGFG36	TTTACAACGT	CGTGACTGGG	AAAACCTGG	CGTTACCCAA	CTTAATCGCC
	3060	3070	3080	3090	3100
1 seq pTGFG36	TTGCAGCACA	TCCCCCTTTC	GCCAGCTGGC	GTAATAGCGA	AGAGGCCCGC
	3110	3120	3130	3140	3150
1 seq pTGFG36	ACCGATCGCC	CTTCCCAACA	GTTGCGCAGC	CTGAATGGCG	AATGGCGCCT
	3160	3170	3180	3190	3200
1 seq pTGFG36	GATGCGGTAT	TTTCTCCTTA	CGCATCTGTG	CGGTATTTC	CACCGCATAT
	3210	3220	3230	3240	3250
1 seq pTGFG36	GGTGCACTCT	CAGTACAATC	TGCTCTGATG	CCGCATAGTT	AAGCCAGCCC
	3260	3270	3280	3290	3300
1 seq pTGFG36	CGACACCCGC	CAACACCCGC	TGACGCGCCC	TGACGGGCTT	GTCTGCTCCC
	3310	3320	3330	3340	3350
1 seq pTGFG36	GGCATCCGCT	TACAGACAAG	CTGTGACCGT	CTCCGGGAGC	TGCATGTGTC
	3360	3370	3380	3390	3400
1 seq pTGFG36	AGAGGTTTTTC	ACCGTCATCA	CCGAAACGCG	CGAGACGAAA	GGGGGGGTAC
	3410	3420	3430	3440	3450
1 seq pTGFG36	CAGCTTCGTA	GCTAGAACAT	CATGTTCTGG	GATATCAGCT	TCGTAGCTAG
	3460	3470	3480	3490	3500
1 seq pTGFG36	AACATCATGT	TCTGGTACCC	CCCTCGTGAT	ACGCCTATTT	TTATAGGTTA
	3510	3520	3530	3540	3550
1 seq pTGFG36	ATGTCATGAT	AATAATGGTT	TCTTAGACGT	CAGGTGGCAC	TTTTCGGGGA
	3560	3570	3580	3590	3600
1 seq pTGFG36	AATGTGCGCG	GAACCCCTAT	TGTTTATTT	TTCTAAATAC	ATTCAAATAT
	3610	3620	3630	3640	3650
1 seq pTGFG36	GTATCCGCTC	ATGAGACAAT	AACCCTGATA	AATGCTTCAA	TAATATTGAA
	3660	3670	3680	3690	3700
1 seq pTGFG36	AAAGGAAGAG	TATGAGTATT	CAACATTTC	GTGTCGCCCT	TATTCCTTT
	3710	3720	3730	3740	3750
1 seq pTGFG36	TTTGCGGCAT	TTGCCTTCC	TGTTTTGCT	CACCCAGAAA	CGCTGGTGAA
	3760	3770	3780	3790	3800
1 seq pTGFG36	AGTAAAAGAT	GCTGAAGATC	AGTTGGGTGC	ACGAGTGGGT	TACATCGAAC

Sequenz pTGFG36 13.12.1998 20:16 Uhr

1000

4

	3810	3820	3830	3840	3850
1 seq pTGFG36	TGGATCTCAA	CAGCGGTAAG	ATCCTTGAGA	GTTTTCGCCC	CGAAGAACGT
	3860	3870	3880	3890	3900
1 seq pTGFG36	TTTCCAATGA	TGAGCACTTT	TAAAGTTCTG	CTATGTGGCG	CGGTATTATC
	3910	3920	3930	3940	3950
1 seq pTGFG36	CCGTATTGAC	GCCGGGCAAG	AGCAACTCGG	TCGCCGCATA	CACTATTCTC
	3960	3970	3980	3990	4000
1 seq pTGFG36	AGAATGACTT	GGTTGAGTAC	TCACCAGTCA	CAGAAAAGCA	TCTTACGGAT
	4010	4020	4030	4040	4050
1 seq pTGFG36	GGCATGACAG	TAAGAGAATT	ATGCAGTGCT	GCCATAACCA	TGAGTGATAA
	4060	4070	4080	4090	4100
1 seq pTGFG36	CACTGCGGCC	AACTTACTTC	TGACAACGAT	CGGAGGACCG	AAGGAGCTAA
	4110	4120	4130	4140	4150
1 seq pTGFG36	CCGCTTTTTT	GCACAACATG	GGGGATCATG	TAAC TCGCCT	TGATCGTTGG
	4160	4170	4180	4190	4200
1 seq pTGFG36	GAACCGGAGC	TGAATGAAGC	CATACCAAAC	GACGAGCGTG	ACACCACGAT
	4210	4220	4230	4240	4250
1 seq pTGFG36	GCCTGTAGCA	ATGGCAACAA	CGTTGCGCAA	ACTATTAAC T	GGCGAACTAC
	4260	4270	4280	4290	4300
1 seq pTGFG36	TTACTCTAGC	TTCCCGCAA	CAATTAATAG	ACTGGATGGA	GGCGGATAAA
	4310	4320	4330	4340	4350
1 seq pTGFG36	GTTGCAGGAC	CACTTCTGCG	CTCGGCCCTT	CCGGCTGGCT	GGTTTATTGC
	4360	4370	4380	4390	4400
1 seq pTGFG36	TGATAAATCT	GGAGCCGGTG	AGCGTG GGTG	TCGCGGTATC	ATTGCAGCAC
	4410	4420	4430	4440	4450
1 seq pTGFG36	TGGGGCCAGA	TGGTAAGCCC	TCCCGTATCG	TAGTTATCTA	CACGACGGGG
	4460	4470	4480	4490	4500
1 seq pTGFG36	AGTCAGGCAA	CTATGGATGA	ACGAAATAGA	CAGATCGCTG	AGATAGGTGC
	4510	4520	4530	4540	4550
1 seq pTGFG36	CTCACTGATT	AAGCATTGGT	AACTGTCAGA	CCAAGTTTAC	TCATATATAC
	4560	4570	4580	4590	4600
1 seq pTGFG36	TTTAGATTGA	TTTAAAACTT	CATTTT TAAT	TTAAAAGGAT	CTAGGTGAAG
	4610	4620	4630	4640	4650
1 seq pTGFG36	ATCCTTTTTG	ATAATCTCAT	GACCAAAATC	CCTTAACGTG	AGTTTTCGTT
	4660	4670	4680	4690	4700
1 seq pTGFG36	CCACTGAGCG	TCAGACCCCG	TAGAAAAGAT	CAAAGGATCT	TCTTGAGATC
	4710	4720	4730	4740	4750
1 seq pTGFG36	CTTTTTTTCT	GCGCGTAATC	TGCTGCTTGC	AAACAAAAAA	ACCACCGCTA

Sequenz pTGFG36 13.12.1998 20:16 Uhr

1 seq pTGFG36	4760	4770	4780	4790	4800
	CCAGCGGTGG	TTGTGTGCC	GGATCAAGAG	CTACCAACTC	TTTTTCCGAA
1 seq pTGFG36	4810	4820	4830	4840	4850
	GGTAACTGGC	TTCAGCAGAG	CGCAGATACC	AAATACTGTT	CTTCTAGTGT
1 seq pTGFG36	4860	4870	4880	4890	4900
	AGCCCGTAGTT	AGGCCACCAC	TTCAAGAACT	CTGTAGCACC	GCCTACATAC
1 seq pTGFG36	4910	4920	4930	4940	4950
	CTCGCTCTGC	TAATCCTGTT	ACCAGTGGCT	GCTGCCAGTG	GCGATAAGTC
1 seq pTGFG36	4960	4970	4980	4990	5000
	GTGTCTTACC	GGGTGGACT	CAAGACGATA	GTTACCGGAT	AAGGCGCAGC
1 seq pTGFG36	5010	5020	5030	5040	5050
	GGTCGGGCTG	AACGGGGGGT	TCGTGCACAC	AGCCCAGCTT	GGAGCGAACG
1 seq pTGFG36	5060	5070	5080	5090	5100
	ACCTACACCG	AACTGAGATA	CCTACAGCGT	GAGCTATGAG	AAAGCGCCAC
1 seq pTGFG36	5110	5120	5130	5140	5150
	GCTTCCCGAA	GGGAGAAAGG	CGGACAGGTA	TCCGGTAAGC	GGCAGGGTCG
1 seq pTGFG36	5160	5170	5180	5190	5200
	GAACAGGAGA	GCGCACGAGG	GAGCTTCCAG	GGGGAAACGC	CTGGTATCTT
1 seq pTGFG36	5210	5220	5230	5240	5250
	TATAGTCCTG	TCGGGTTTCG	CCACCTCTGA	CTTGAGCGTC	GATTTTTGTG
1 seq pTGFG36	5260	5270	5280	5290	5300
	ATGCTCGTCA	GGGGGGCGGA	GCCTATGGAA	AAACGCCAGC	AACGCGGCCT
1 seq pTGFG36	5310	5320	5330	5340	5350
	TTTTACGGTT	CCTGGCCTTT	TGCTGGCCTT	TTGCTCACAT	GTTCTTTCCT
1 seq pTGFG36	5360	5370	5380	5390	5400
	GCGTTATCCC	CTGATTCTGT	GGATAACCGT	ATTACCGCCT	TTGAGTGAGC
1 seq pTGFG36	5410	5420	5430	5440	5450
	TGATACCGCT	CGCCGCAGCC	GAACGACCGA	GCGCAGCGAG	TCAGTGAGCG
1 seq pTGFG36	5460	5470	5480	5490	5500
	AGGAAGCGGA	AGAGCGCCCA	ATACGCAAAC	CGCCTCTCCC	CGCGCGTTGG
1 seq pTGFG36	5510	5520	5530	5540	5550
	CCGATTCAAT	AATGCAGCTG	GCACGACAGG	TTCCCGACT	GGAAAGCGGG
1 seq pTGFG36	5560	5570	5580	5590	5600
	CAGTGAGCGC	AACGCAATTA	ATGTGAGTTA	GCTCACTCAT	TAGGCACCCC
1 seq pTGFG36	5610	5620	5630	5640	5650
	AGGCTTTACA	CTTTATGCTT	CCGGCTCGTA	TGTGTGTGG	AATTGTGAGC
1 seq pTGFG36	5660	5670	5680	5690	5700
	GGATAACAAT	TTCACACAGG	AAACAGCTAT	GACCATGATT	ACGCCAAGCT

Sequenz pTGFG36 13.12.1998 20:16 Uhr

10000

	5710	5720	5730	5740	5750
1 seq pTGFG36	CTCTAGAGCT	CTAGAGCTCT	AGAGCTCTAG	AGAGCTTGCA	TGCCTGCAGG
	5760	5770	5780	5790	5800
1 seq pTGFG36	TCG				

sequenz pTGFG67 13.12.1998 21:02 Uhr

	0	20	40	60	80	100
1 seq pTGFG67	CGCGTTGACA	TTGATTATTG	ACTAGTTATT	AATAGTAATC	AATTACGGGG	
	60	70	80	90	100	
1 seq pTGFG67	TCATTAGTTC	ATAGCCCATATA	TATGGAGTTC	CGCGTTACAT	AACTTACGGT	
	110	120	130	140	150	
1 seq pTGFG67	AAATGGCCCCG	CCTGGCTGAC	CGCCCAACGA	CCCCCGCCCA	TTGACGTCAA	
	160	170	180	190	200	
1 seq pTGFG67	TAATGACGTA	TGTTCCCATATA	GTAACGCCAA	TAGGGACTTT	CCATTGACGT	
	210	220	230	240	250	
1 seq pTGFG67	CAATGGGTGG	AGTATTTACG	GTAAACTGCC	CACTTGGCAG	TACATCAAGT	
	260	270	280	290	300	
1 seq pTGFG67	GTATCATATG	CCAAGTACGC	CCCCTATTGA	CGTCAATGAC	GGTAAATGGC	
	310	320	330	340	350	
1 seq pTGFG67	CCGCCTGGCA	TTATGCCCAG	TACATGACCT	TATGGGACTT	TCCTACTTGG	
	360	370	380	390	400	
1 seq pTGFG67	CAGTACATCT	ACGTATTAGT	CATCGCTATT	ACCATGGTGA	TGCGGTTTTC	
	410	420	430	440	450	
1 seq pTGFG67	GCAGTACATC	AATGGGCGTG	GATAGCGGTT	TGACTCACGG	GGATTTCCAA	
	460	470	480	490	500	
1 seq pTGFG67	GTCTCCACCC	CATTGACGTC	AATGGGAGTT	TGTTTTGGCA	CCAAAATCAA	
	510	520	530	540	550	
1 seq pTGFG67	CGGGACTTTC	CAAAATGTCG	TAACAACTCC	GCCCCATTGA	CGCAAATGGG	
	560	570	580	590	600	
1 seq pTGFG67	CGGTAGGCGT	GTACGGTGGG	AGGTCTATAT	AAGCAGAGCT	CTCTGGCTAA	
	610	620	630	640	650	
1 seq pTGFG67	CTAGAGAACC	CACTGCTTAC	TGGCTTATCG	AAATTAATAC	GACTCACTAT	
	660	670	680	690	700	
1 seq pTGFG67	AGGGAGACCC	AAGCTTGACC	TCGAGCAAGC	GGCCGCGACT	CTACTAGAGG	
	710	720	730	740	750	
1 seq pTGFG67	ATCTTTGTGA	AGGAACCTTA	CTTCTGTGGT	GTGACATAAT	TGGACAAACT	
	760	770	780	790	800	
1 seq pTGFG67	ACCTACAGAG	ATTAAAGCT	CTAAGGTAAA	TATAAAATTT	TTAAGTGTAT	
	810	820	830	840	850	
1 seq pTGFG67	AATGTGTTAA	ACTACTGATT	CTAATGTTT	GTGTATTTTA	GATTCCAACC	
	860	870	880	890	900	
1 seq pTGFG67	TATGGAAGT	ATGAATGGGA	GCAGTGGTGG	AATGCCTTTA	ATGAGGAAAA	
	910	920	930	940	950	
1 seq pTGFG67	CCTGTTTTGC	TCAGAAGAAA	TGCCATCTAG	TGATGATGAG	GCTACTGCTG	

1000

Sequenz pTGFG67 13.12.1998 21:02 Uhr

	960	970	980	990	1000
1 seq pTGFG67	ACTCTCAACA TTCTACTCCT CCAAAAAAGA AGAGAAAGGT AGAAGACCCC				
	1010	1020	1030	1040	1050
1 seq pTGFG67	AAGGACTTTC CTTCAGAATT GCTAAGTTTT TTGAGTCATG CTGTGTTTAG				
	1060	1070	1080	1090	1100
1 seq pTGFG67	TAATAGAACT CTTGCTTGCT TTGCTATTTA CACCACAAAG GAAAAAGCTG				
	1110	1120	1130	1140	1150
1 seq pTGFG67	CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC CTTTATAAGT				
	1160	1170	1180	1190	1200
1 seq pTGFG67	AGGCATAACA GTTATAATCA TAACATACTG TTTTTTCTTA CTCCACACAG				
	1210	1220	1230	1240	1250
1 seq pTGFG67	GCATAGAGTG TCTGCTATTA ATAACATGCT TCAAAAATTG TGTACCTTTA				
	1260	1270	1280	1290	1300
1 s	GCTTTTTAAT TTGTAAAGGG GTTAATAAGG AATATTTGAT GTATAGTGCC				
	1310	1320	1330	1340	1350
1 seq pTGFG67	TTGACTAGAG ATCATAATCA GCCATACCAC ATTTGTAGAG GTTTTACTTG				
	1360	1370	1380	1390	1400
1 seq pTGFG67	CTTTAAAAAA CCTCCACAC CTCCCCCTGA ACCTGAAACA TAAATGAAT				
	1410	1420	1430	1440	1450
1 seq pTGFG67	GCAATTGTIG TTGTTAACTT GTTTATTGCA GCTTATAATG GTTACAAATA				
	1460	1470	1480	1490	1500
1 seq pTGFG67	AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTTTT TCACTGCATT				
	1510	1520	1530	1540	1550
1 seq pTGFG67	CTAGTTGTGG TTGTCCAAA CTCATCAATG TATCTTATCA TGTCTGGATC				
	1560	1570	1580	1590	1600
1 seq pTGFG67	CCCGGGTACC CTCTAGAGCG AATTAATTCA CTGGCCGTCG TTTTACAACG				
	1610	1620	1630	1640	1650
1 seq pTGFG67	TCGTGACTGG GAAAACCCTG GCGTTACCCA ACTTAATCGC CTGTCAGCAC				
	1660	1670	1680	1690	1700
1 seq pTGFG67	ATCCCCCTTT CGCCAGCTGG CGTAATAGCG AAGAGGCCCG CACCGATCGC				
	1710	1720	1730	1740	1750
1 seq pTGFG67	CCTTCCCAAC AGTTGCGCAG CCTGAATGGC GAATGGCGCC TGATGCGGTA				
	1760	1770	1780	1790	1800
1 seq pTGFG67	TTTTCTCCTT ACGCATCTGT GCGGTATTTT ACACCGCATA TGGTGCATCT				
	1810	1820	1830	1840	1850
1 seq pTGFG67	TCAGTACAAT CTGCTCTGAT GCCGCATAGT TAAGCCAGCC CCGACACCCG				
	1860	1870	1880	1890	1900
1 seq pTGFG67	CCAACACCCG CTGACGCGCC CTGACGGGCT TGTCTGCTCC CGGCATCCGC				

Sequenz pTGFG67 13.1.1998 21:02 Uhr

	1910	1920	1930	1940	1950
1 seq pTGFG67	TTACAGACAA	GCTGTGACCG	TCTCCGGGAG	CTGCATGTGT	CAGAGGTTTT
	1960	1970	1980	1990	2000
1 seq pTGFG67	CACCGTCATC	ACCGAAACGC	GCGAGACGAA	AGGGGGGGTA	CCAGCTTCGT
	2010	2020	2030	2040	2050
1 seq pTGFG67	AGCTAGAACA	TCATGTTCTG	GGATATCAGC	TTCGTAGCTA	GAACATCATG
	2060	2070	2080	2090	2100
1 seq pTGFG67	TTCTGGTACC	CCCCTCGTGA	TACGCCTATT	TTTATAGGTT	AATGTCATGA
	2110	2120	2130	2140	2150
1 seq pTGFG67	TAATAATGGT	TTCTTAGACG	TCAGGTGGCA	CTTTTCGGGG	AAATGTGCGC
	2160	2170	2180	2190	2200
1 seq pTGFG67	GGAACCCCTA	TTGTTTATT	TTTCTAAATA	CATTCAAATA	TGTATCCGCT
	2210	2220	2230	2240	2250
1 seq pTGFG67	CATGAGACAA	TAACCCTGAT	AAATGCTTCA	ATAATATTGA	AAAAGGAAGA
	2260	2270	2280	2290	2300
1 seq pTGFG67	GTATGAGTAT	TCAACATTTC	CGTGTCGCCC	TTATTCCCTT	TTTGCGGCA
	2310	2320	2330	2340	2350
1 seq pTGFG67	TTTGCCCTTC	CTGTTTTTGC	TCACCCAGAA	ACGCTGGTGA	AAGTAAAAGA
	2360	2370	2380	2390	2400
1 seq pTGFG67	TGCTGAAGAT	CAGTTGGGTG	CACGAGTGGG	TTACATCGAA	CTGGATCTCA
	2410	2420	2430	2440	2450
1 seq pTGFG67	ACAGCGGTAA	GATCCTTGAG	AGTTTTCGCC	CCGAAGAACG	TTTTCCAATG
	2460	2470	2480	2490	2500
1 seq pTGFG67	ATGAGCACTT	TTAAAGTTCT	GCTATGTGGC	GCGGTATTAT	CCCGTATTGA
	2510	2520	2530	2540	2550
1 seq pTGFG67	CGCCGGGCAA	GAGCAACTCG	GTCGCCGCAT	ACACTATTCT	CAGAATGACT
	2560	2570	2580	2590	2600
1 seq pTGFG67	TGGTTGAGTA	CTCACCAGTC	ACAGAAAAGC	ATCTTACGGA	TGGCATGACA
	2610	2620	2630	2640	2650
1 seq pTGFG67	GTAAGAGAAT	TATGCAGTGC	TGCCATAACC	ATGAGTGATA	ACACTGCGGC
	2660	2670	2680	2690	2700
1 seq pTGFG67	CAACTTACTT	CTGACAACGA	TCGGAGGACC	GAAGGAGCTA	ACCGCTTTTT
	2710	2720	2730	2740	2750
1 seq pTGFG67	TGCACAACAT	GGGGGATCAT	GTAACGCGCC	TTGATCGTTG	GGAACCGGAG
	2760	2770	2780	2790	2800
1 seq pTGFG67	CTGAATGAAG	CCATACCAA	CGACGAGCGT	GACACCACGA	TGCCTGTAGC
	2810	2820	2830	2840	2850
1 seq pTGFG67	AATGGCAACA	ACGTTGCGCA	AACTATTAAC	TGGCGAACTA	CTTACTCTAG


Sequenz pTGFG67 13.12.1998 21:02 Uhr

	2850	2870	2880	2890	2900
1 seq pTGFG67	CTTCCCGGCA ACAATTAATA GACTGGATGG AGGCGGATAA AGTTGCAGGA				
	2910	2920	2930	2940	2950
1 seq pTGFG67	CCACTTCTGC GCTCGGCCCT TCCGGCTGGC TGGTTTATTG CTGATAAATC				
	2960	2970	2980	2990	3000
1 seq pTGFG67	TGGAGCCGGT GAGCGTGGGT CTCGCGGTAT CATTGCAGCA CTGGGGCCAG				
	3010	3020	3030	3040	3050
1 seq pTGFG67	ATGGTAAGCC CTCCCGTATC GTAGTTATCT ACACGACGGG GAGTCAGGCA				
	3060	3070	3080	3090	3100
1 seq pTGFG67	ACTATGGATG AACGAAATAG ACAGATCGCT GAGATAGGTG CCTCACTGAT				
	3110	3120	3130	3140	3150
1 seq pTGFG67	TAAGCATTGG TAACTGTCAG ACCAAGTTTA CTCATATATA CTTTAGATTG				
	3160	3170	3180	3190	3200
1 seq pTGFG67	ATTTAAAACT TCATTTTAA TTTAAAAGGA TCTAGGTGAA GATCCTTTTT				
	3210	3220	3230	3240	3250
1 seq pTGFG67	GATAATCTCA TGACCAAAT CCCTTAACGT GAGTTTTCGT TCCACTGAGC				
	3260	3270	3280	3290	3300
1 seq pTGFG67	GTCAGACCCC GTAGAAAAGA TCAAAGGATC TTCTTGAGAT CCTTTTTTTC				
	3310	3320	3330	3340	3350
1 seq pTGFG67	TGCGCGTAAT CTGCTGCTTG CAAACAAAAA AACCACCGCT ACCAGCGGTG				
	3360	3370	3380	3390	3400
1 seq pTGFG67	GTTTGTTTGC CGGATCAAGA GCTACCAACT CTTTTTCCGA AGGTAAGTGG				
	3410	3420	3430	3440	3450
1 seq pTGFG67	CTTCAGCAGA GCGCAGATAC CAAATACTGT TCTTCTAGTG TAGCCGTAGT				
	3460	3470	3480	3490	3500
1 seq pTGFG67	TAGGCCACCA CTCAAGAAC TCTGTAGCAC CGCCTACATA CCTCGCTCTG				
	3510	3520	3530	3540	3550
1 seq pTGFG67	CTAATCCTGT TACCAGTGGC TGCTGCCAGT GCGATAAGT CGTGTCTTAC				
	3560	3570	3580	3590	3600
1 seq pTGFG67	CGGGTTGGAC TCAAGACGAT AGTTACCGGA TAAGGCGCAG CGGTCGGGCT				
	3610	3620	3630	3640	3650
1 seq pTGFG67	GAACGGGGGG TTCGTGCACA CAGCCAGCT TGGAGCGAAC GACCTACACC				
	3660	3670	3680	3690	3700
1 seq pTGFG67	GAACTGAGAT ACCTACAGCG TGAGCTATGA GAAAGCGCCA CGCTTCCCGA				
	3710	3720	3730	3740	3750
1 seq pTGFG67	AGGGAGAAAG GCGGACAGGT ATCCGGTAAG CGGCAGGGTC GGAACAGGAG				
	3760	3770	3780	3790	3800
1 seq pTGFG67	AGCGCACGAG GGAGCTTCCA GGGGAAACG CCTGGTATCT TTATAGTCCT				

Sequenz pTGFG67 13.12.98 21:02 Uhr

	3810	3820	3830	3840	3850
1 seq pTGFG67	GTCGGGTTTC	GCCACCTCTG	ACTTGAGCGT	CGATTTTGT	GATGCTCGTC
	3860	3870	3880	3890	3900
1 seq pTGFG67	AGGGGGGCGG	AGCCTATGGA	AAAACGCCAG	CAACGCGGCC	TTTTTACGGT
	3910	3920	3930	3940	3950
1 seq pTGFG67	TCCTGGCCTT	TTGCTGGCCT	TTTGCTCACA	TGTTCTTTCC	TGCGTTATCC
	3960	3970	3980	3990	4000
1 seq pTGFG67	CCTGATTCTG	TGGATAACCG	TATTACCGCC	TTTGAGTGAG	CTGATACCGC
	4010	4020	4030	4040	4050
1 seq pTGFG67	TCGCCGCAGC	CGAACGACCG	AGCGCAGCGA	GTCAGTGAGC	GAGGAAGCGG
	4060	4070	4080	4090	4100
1 seq pTGFG67	AAGAGCGCCC	AATACGCAAA	CCGCCTCTCC	CCGCGCGTTG	GCCGATTCAT
	4110	4120	4130	4140	4150
1 seq pTGFG67	TAATGCAGCT	GGCACGACAG	GTTTCCCGAC	TGGAAAGCGG	GCAGTGAGCG
	4160	4170	4180	4190	4200
1 seq pTGFG67	CAACGCAATT	AATGTGAGTT	AGCTCACTCA	TTAGGCACCC	CAGGCTTTAC
	4210	4220	4230	4240	4250
1 seq pTGFG67	ACTTTATGCT	TCCGGCTCGT	ATGTTGTGTG	GAATTGTGAG	CGGATAACAA
	4260	4270	4280	4290	4300
1 seq pTGFG67	TTTCACACAG	GAAACAGCTA	TGACCATGAT	TACGCCAAGC	TCTCTAGAGC
	4310	4320	4330	4340	4350
1 seq pTGFG67	TCTAGAGCTC	TAGAGCTCTA	GAGAGCTTGC	ATGCCTGCAG	GTCG

Figure 47



Concentration of Expressed Marker Gene "Green Fluorescent Protein" (GFP)
versus "Bovine Serum Albumin" (BSA)



Figure 48



Figure 48a




Figure 48b




Figure 48c




Figure 48d

Corresponding light (a and c) and fluorescent (b and d) microscopic views of HeLa cells transfected with pTGFG5 (a and b) and pTGFG20 (c and d) respectively.

THIS PAGE BLANK (USPTO)